

Lampiran 1. Kuisisioner Yang Akan Dibagikan Kepada Responden

1. Nama lengkap :
2. Jenis Kelamin : Pria / Wanita
3. Umur : Tahun
4. Status : Kawin/ Belum Kawin
5. Pendidikan Terakhir :

Petunjuk pengisian Kuesioner :

1. Bacalah setiap pertanyaan dengan baik dan teliti.
2. Pilih salah satu jawaban yang paling tepat menurut Bapak/ Ibu.

Yaitu dengan cara memberi tanda *check list* (√) pada kolom jawaban yang tersedia.

Keterangan Jawaban Kuesioner :

- | | |
|-----------------------------|-----------------|
| 1.Sangat Tidak Setuju (STS) | bobot nilai = 1 |
| 2.TidakSetuju (TS) | bobot nilai = 2 |
| 3.Netral (N) | bobot nilai = 3 |
| 4.Setuju (S) | bobot nilai = 4 |
| 5.Sangat Setuju (SS) | bobot nilai = 5 |

		Variabel : <i>Game Online</i> (X₁)				
No	Pernyataan	Jawaban				
		SS	S	N	TS	STS
X _{1.1}	Saya sering bermain <i>game online</i> di rumah dan sekolah					
	Saya sering bermain <i>game online</i> di sembarang tempat					

X.1.2	Saya sering bermain <i>game online</i> saat orang tua tidak ada dirumah					
	Saya sering bermain <i>game online</i> saat waktu libur sekolah					
X.1.3	Saya sering bermain <i>game online</i> yang berbayar					
	Saya sering bermain <i>game online</i> yang gratisan					
X.1.4	Saya lebih memilih bermain <i>game online</i> dulu dari pada mengerjakan PR					
	Saya sering bermain <i>game online</i> dari pada masuk sekolah					
Variabel : Perilaku Remaja (Y₁)						
Y.1.1	saya ketika bermain <i>game online</i> sering melupakan makan					
	Saya ketika bermain <i>game online</i> sering meninggalkan les saya					
Y.1.2	Saya merasa dari bermain <i>game online</i> saya semakin sering tidak membantu					

	pekerjaan orang tua saya					
	Saya merasa karena keseringan bermain <i>game online</i> saya jarang mempunyai teman					

Lampiran 2. Data Tabulasi

X1.1	X1.A	X1.2	X1.A	X1.3	X1.A	X1.4	X1.A	Y1.1	Y1.A	Y1.2	Y1.A
4	5	5	5	4	5	4	5	4	5	5	5
4	5	5	5	5	4	4	5	4	4	5	4
4	4	4	4	4	3	4	5	4	4	4	4
5	5	5	5	4	5	5	4	5	4	3	3
5	5	5	4	5	5	4	5	5	4	5	5
4	5	5	4	5	4	5	4	4	4	4	4
4	5	5	5	3	5	5	5	5	5	5	5
4	4	4	3	4	4	4	4	4	4	4	4
4	4	5	5	4	4	4	4	4	4	4	4
4	4	4	4	3	4	4	3	4	4	3	4
4	4	4	2	3	4	3	4	4	2	4	4
4	5	4	3	4	4	4	4	4	3	4	4
5	4	4	5	5	4	4	4	4	4	4	4
5	4	4	5	5	4	4	4	4	4	4	4
4	4	4	3	2	4	4	5	1	3	4	4
4	4	4	5	3	4	3	2	4	4	2	4
4	4	3	3	4	4	4	4	4	4	4	4
4	4	4	3	3	3	3	3	4	3	4	4
4	4	4	3	4	4	3	4	3	4	4	3
4	4	4	4	3	4	4	3	3	4	3	3
4	4	3	4	4	3	3	3	3	4	4	4
4	4	3	4	4	4	4	4	4	4	4	4
4	4	3	4	4	3	4	3	4	3	4	3

4	4	4	4	3	4	4	4	3	3	4	4
4	3	4	4	3	3	3	3	3	3	4	3
4	4	3	1	1	3	3	3	1	1	1	3
4	4	4	3	4	4	5	3	2	3	5	4
4	4	4	4	5	4	4	3	4	4	4	4
4	5	5	4	4	5	5	4	4	4	4	5
4	3	4	4	3	3	3	3	3	3	4	3

Lampiran 3. Hasil Hitung SPSS 22.0

Notes		
Output Created		27-JUL-2020 07:55:22
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X1.1 X1.1_A X1.2 X1.2_A X1.3 X1.3_A X1.4 X1.4_A Y1.1 Y1.1_A Y1.2 Y1.2_A /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.05
	Elapsed Time	00:00:00.08

Correlations

Notes		
Output Created		27-JUL-2020 07:56:32
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30

X1.4_A	Pearson Correlation	.223	.560**	.492**	.185	.290	.509**	.414*	1	.258	.360	.578**	.522**	.646**
	Sig. (2-tailed)	.236	.001	.006	.328	.120	.004	.023		.169	.051	.001	.003	.000
	N	30	30	30	30	30	30	30	30	30	30	30	30	30
Y1.1	Pearson Correlation	.359	.467**	.438*	.571**	.608**	.462*	.319	.258	1	.618**	.334	.406*	.749**
	Sig. (2-tailed)	.052	.009	.016	.001	.000	.010	.086	.169		.000	.071	.026	.000
	N	30	30	30	30	30	30	30	30	30	30	30	30	30
Y1.1_A	Pearson Correlation	.181	.403*	.457*	.768**	.590**	.551**	.445*	.360	.618**	1	.449*	.484**	.799**
	Sig. (2-tailed)	.339	.027	.011	.000	.001	.002	.014	.051	.000		.013	.007	.000
	N	30	30	30	30	30	30	30	30	30	30	30	30	30
Y1.2	Pearson Correlation	.047	.267	.389*	.319	.570**	.306	.352	.578**	.334	.449*	1	.517**	.665**
	Sig. (2-tailed)	.804	.154	.034	.086	.001	.100	.057	.001	.071	.013		.003	.000
	N	30	30	30	30	30	30	30	30	30	30	30	30	30
Y1.2_A	Pearson Correlation	.066	.577**	.455*	.268	.313	.603**	.403*	.522**	.406*	.484**	.517**	1	.722**
	Sig. (2-tailed)	.730	.001	.012	.152	.093	.000	.027	.003	.026	.007	.003		.000
	N	30	30	30	30	30	30	30	30	30	30	30	30	30
SKOR	Pearson Correlation	.372*	.703**	.703**	.670**	.693**	.740**	.621**	.646**	.749**	.799**	.665**	.722**	1
	Sig. (2-tailed)	.043	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	
	N	30	30	30	30	30	30	30	30	30	30	30	30	30

*. Correlation is significant at the 0.05 level (2-tailed).

** . Correlation is significant at the 0.01 level (2-tailed).

RELIABILITY

```

/VARIABLES=X1.1 X1.1_A X1.2 X1.2_A X1.3 X1.3_A X1.4 X1.4_A Y1.1 Y1.1_A Y1.2 Y1.2_A SKOR
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA.

```


Reliability

Notes		
Output Created		27-JUL-2020 07:57:09
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Matrix Input	
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.
Syntax		RELIABILITY /VARIABLES=X1.1 X1.1_A X1.2 X1.2_A X1.3 X1.3_A X1.4 X1.4_A Y1.1 Y1.1_A Y1.2 Y1.2_A SKOR /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA.
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.01

Scale: ALL VARIABLES

Case Processing Summary			
		N	%
Cases	Valid	30	100.0
	Excluded ^a	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

Reliability Statistics

Cronbach's Alpha	N of Items
.096	13

NPAR TESTS

```
/K-S (NORMAL)=X1.1 X1.1_A X1.2 X1.2_A X1.3 X1.3_A X1.4 X1.4_A Y1.1 Y1.1_A Y1.2 Y1.2_A
/MISSING ANALYSIS.
```

NPar Tests

Notes

Output Created	27-JUL-2020 07:57:45	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each test are based on all cases with valid data for the variable(s) used in that test.
Syntax	NPAR TESTS /K-S(NORMAL)=X1.1 X1.1_A X1.2 X1.2_A X1.3 X1.3_A X1.4 X1.4_A Y1.1 Y1.1_A Y1.2 Y1.2_A /MISSING ANALYSIS.	
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.04
	Number of Cases Allowed ^a	52428

a. Based on availability of workspace memory.

One-Sample Kolmogorov-Smirnov Test

		X1.1	X1.1_A	X1.2	X1.2_A	X1.3	X1.3_A	X1.4	X1.4_A	Y1.1	Y1.1_A	Y1.2	Y1.2_A
N		30	30	30	30	30	30	30	30	30	30	30	30
Normal Parameters ^{a,b}	Mean	4.1333	4.2000	4.1000	3.8667	3.7333	3.9333	3.9000	3.8000	3.6333	3.6333	3.9000	3.9000
	Std. Deviation	.34575	.55086	.66176	.97320	.94443	.63968	.66176	.80516	.96431	.80872	.84486	.60743
Most Extreme Differences	Absolute	.517	.375	.293	.254	.244	.308	.293	.231	.348	.342	.380	.332
	Positive	.517	.375	.293	.179	.189	.292	.273	.206	.252	.258	.286	.301
	Negative	-.350	-.292	-.273	-.254	-.244	-.308	-.293	-.231	-.348	-.342	-.380	-.332
Test Statistic		.517	.375	.293	.254	.244	.308	.293	.231	.348	.342	.380	.332
Asymp. Sig. (2-tailed)		.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c	.000 ^c

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

Regression

Notes

Output Created		27-JUL-2020 08:00:02
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any variable used.

Syntax	REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA COLLIN TOL /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT Y /METHOD=ENTER X /RESIDUALS HISTOGRAM(ZRESID) NORMPROB(ZRESID) /SAVE RESID.	
Resources	Processor Time	00:00:02.37
	Elapsed Time	00:00:02.32
	Memory Required	1620 bytes
	Additional Memory Required for Residual Plots	656 bytes
Variables Created or Modified	RES_1	Unstandardized Residual

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	X ^b	.	Enter

a. Dependent Variable: Y

b. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.806 ^a	.649	.637	1.50922

a. Predictors: (Constant), X

b. Dependent Variable: Y

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	118.090	1	118.090	51.845	.000 ^b
	Residual	63.777	28	2.278		
	Total	181.867	29			

a. Dependent Variable: Y

b. Predictors: (Constant), X

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.	Collinearity Statistics	
		B	Std. Error	Beta			Tolerance	VIF
1	(Constant)	-1.216	2.278		-.534	.598		
	X	.514	.071	.806	7.200	.000	1.000	1.000

a. Dependent Variable: Y

Collinearity Diagnostics^a

Model	Dimension	Eigenvalue	Condition Index	Variance Proportions	
				(Constant)	X
1	1	1.993	1.000	.00	.00
	2	.007	16.474	1.00	1.00


a. Dependent Variable: Y

Residuals Statistics^a


	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	10.0963	18.3231	15.0667	2.01794	30
Residual	-4.09627	2.33284	.00000	1.48297	30
Std. Predicted Value	-2.463	1.614	.000	1.000	30
Std. Residual	-2.714	1.546	.000	.983	30

a. Dependent Variable: Y

Charts



Normal P-P Plot of Regression Standardized Residual



Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item- Total Correlation	Cronbach's Alpha if Item Deleted
X1.1	42.6000	35.903	.360	.893
X1.1_A	42.5333	33.085	.651	.882
X1.2	42.6333	32.102	.663	.880
X1.2_A	42.8667	30.120	.603	.884
X1.3	43.0000	30.000	.640	.881
X1.3_A	42.8000	32.028	.701	.878
X1.4	42.8333	32.557	.598	.883
X1.4_A	42.9333	31.857	.551	.885
Y1.1	43.1000	29.748	.650	.881
Y1.1_A	43.1000	30.231	.745	.874
Y1.2	42.8333	31.454	.564	.885
Y1.2_A	42.8333	32.902	.609	.883